

# Ventilatory Response to Nitrogen Multiple-Breath Washout in Infants

Florian Singer, MD, PhD,<sup>1,2\*</sup> Sophie Yammine, MD,<sup>1</sup> Anne Schmidt, MD,<sup>1,3</sup> Elena Proietti, MD,<sup>1,3</sup> Elisabeth Kieninger, MD, PhD,<sup>1</sup> Juerg Barben, MD,<sup>4</sup> Carmen Casaulta, MD,<sup>1</sup> Nicolas Regamey, MD,<sup>1</sup> Per Gustafsson, MD, PhD,<sup>5</sup> Urs Frey, MD, PhD,<sup>3</sup> and Philipp Latzin, MD, PhD<sup>1</sup>

**Summary.** Background: Nitrogen multiple-breath washout (N<sub>2</sub>MBW) using 100% oxygen (O<sub>2</sub>) has regained interest to assess efficiency of tracer gas clearance in, for example, children with Cystic Fibrosis (CF). However, the influence of hyperoxia on the infants' respiratory control is unclear. We assessed safety and impact on breathing pattern from hyperoxia, and if exposure to 40% O<sub>2</sub> first induces tolerance to subsequent 100% O<sub>2</sub> for N<sub>2</sub>MBW. Methods: We prospectively enrolled 39 infants aged 3–57 weeks: 15 infants with CF (8 sedated for testing) and 24 healthy controls. Infants were consecutively allocated to the protocols comprising of 100% O<sub>2</sub> or 40/100% O<sub>2</sub> administered for 30 breaths. Lung function was measured using an ultrasonic flowmeter setup. Primary outcome was tidal volume (V<sub>T</sub>). Results: None of the infants experienced apnea, desaturation, or bradycardia. Both protocols initially induced hypoventilation. V<sub>T</sub> temporarily declined in 33/39 infants across 10–25 breaths. Hypoventilation occurred independent of age, disease, and sedation. In the new 40/100% O<sub>2</sub> protocol, V<sub>T</sub> returned to baseline during 40% O<sub>2</sub> and remained stable during 100% O<sub>2</sub> exposure. End-tidal carbon dioxide monitored online did not change. Conclusion: The classical N<sub>2</sub>MBW protocol with 100% O<sub>2</sub> may change breathing patterns of the infants. The new protocol with 40% O<sub>2</sub> induces hyperoxia-tolerance and does not lead to changes in breathing patterns during later N<sub>2</sub> washout using 100% O<sub>2</sub>. Both protocols are safe, the new protocol seems an attractive option for N<sub>2</sub>MBW in infants.

**Pediatr Pulmonol.** © 2013 Wiley Periodicals, Inc.

**Key words:** cystic fibrosis; inert gases; lung function tests; respiratory physiology.

**Funding source:** none reported

## BACKGROUND

Multiple-breath inert tracer gas washout (MBW) is an easy and sensitive lung function test assessing efficiency of inert tracer gas clearance across multiple tidal breaths.<sup>1–4</sup> In children with cystic fibrosis (CF), impaired tracer gas clearance quantified by the lung clearance index (LCI) correlates to other biomarkers of small airways disease,<sup>5–8</sup> and detects pulmonary function response upon short-term interventions.<sup>9–11</sup>

To overcome the need for expensive inert tracer gases such as sulfur hexafluoride, nitrogen (N<sub>2</sub>) MBW using

pure oxygen (O<sub>2</sub>) has regained interest and seems attractive for MBW in infants. However, there is scanty evidence as to whether 100% O<sub>2</sub> for the purpose of MBW is applicable in infants.<sup>12–14</sup> A regular breathing pattern is a prerequisite for reliable MBW testing. By unloading peripheral chemoreceptors, hyperoxia may change tidal volume (V<sub>T</sub>) if 100% O<sub>2</sub> is used.<sup>15</sup> We hypothesized that 40% O<sub>2</sub> may induce “tolerance” of the infants' respiratory control to subsequent 100% O<sub>2</sub> exposure N<sub>2</sub>MBW. Breathing 40% O<sub>2</sub> would allow later washout of the lungs' remaining N<sub>2</sub> fraction using pure O<sub>2</sub>. In the current study, we measured tidal breathing indices when infants

<sup>1</sup>Department of Paediatrics, University Hospital of Bern, Bern, Switzerland.

<sup>2</sup>University Children's Hospital of Zurich, Zurich, Switzerland.

<sup>3</sup>University Children's Hospital of Basel, Basel, Switzerland.

<sup>4</sup>Paediatric Pulmonology, Children's Hospital, St. Gallen, Switzerland.

<sup>5</sup>Department of Paediatrics, Central Hospital, Skoevde, Sweden.

\*Correspondence to: Florian Singer, MD, PhD, University Children's Hospital of Zurich, Zurich 8032, Switzerland.  
E-mail: florian.singer@kispi.uzh.ch

Received 31 March 2013; Accepted 31 May 2013.

DOI 10.1002/ppul.22841  
Published online in Wiley Online Library  
(wileyonlinelibrary.com).

Conflict of interest: None.

© 2013 Wiley Periodicals, Inc.

were exposed to hyperoxia during the classical 1-step N<sub>2</sub>MBW protocol with a switch from ambient to 100% O<sub>2</sub> gas, and the new 2-steps protocol introducing an intermediate 40% O<sub>2</sub> step, that is, 40/100% O<sub>2</sub>. Our aim was to assess safety and impact of hyperoxia (40% or 100% O<sub>2</sub>) on breathing patterns in infants with CF and healthy controls, and to compare tidal breathing parameters from both N<sub>2</sub>MBW protocols. Primary outcome was change in V<sub>T</sub> upon 40% or 100% O<sub>2</sub> exposure.

## METHODS

We prospectively enrolled 40 infants (16 with CF) from two ongoing studies, the *Bern Infant Lung Development (BILD)* cohort and the *Swiss CF Newborn Screening Study*.<sup>16–18</sup> The following inclusion criteria applied: white ethnicity, term delivery (37 weeks), and no known other diseases likely to affect lung function. General exclusion criteria were clinical signs of lower respiratory tract infection. The population is described in Table 1. The study was approved by the Ethics Committee of the Canton of Bern, Bern, Switzerland. Parents provided written informed consent for this study. Parents from one child withdrew consent.

We consecutively allocated 23 infants (7 with CF of which 4 were sedated) to protocol I and 16 (8 with CF of which 4 were sedated) to protocol II. The sequence of tidal breathing measurements in protocol I was (i) 1 baseline test during ambient air exposure (21% O<sub>2</sub>) and (ii) 1 test during breathing 100% O<sub>2</sub>. Protocol II implied (i) 1 baseline test during ambient air exposure, (ii) 1 test during breathing 40% O<sub>2</sub>, and (iii) 1 test during breathing 100% O<sub>2</sub>. Hyperoxia was administered for 30 breaths which reflect approximately twice the cumulative expired volume (CEV) of average N<sub>2</sub>MBW in infants.<sup>12</sup> Oxygen saturation, heart rate, and the exhaled carbon dioxide (CO<sub>2</sub>) fraction were monitored online.

**TABLE 1—Anthropometric Characteristics of the Study Population (n = 39)**

	Younger infants	Older infants
N	31 (7 with CF)	8 (8 with CF)
Protocol I/II	19/12	4/4
Males/females	16/15	3/5
Age	5 (5–6) weeks	13 (13–14) months
Weight	4,150 (3,600–4,960) g	9.3 (8.7–9.5) kg
Height	55 (52–58) cm	77 (73–78) cm

Anthropometric data from the 24 healthy infants and 15 infants with CF. Numeric data are described as median and interquartile range. Protocol I = classical 1-step nitrogen multiple-breath washout protocol using 100% oxygen; Protocol II = new 2-steps nitrogen multiple-breath washout protocol using 40–100% oxygen. Lung function was measured during natural quiet sleep in younger infants and during sedation in older infants.


Lung function was measured in infants in supine position with the head midline via an infant mask (size 1; Homedica AG, Huenenberg, Switzerland), according to the ERS/ATS standards of infant lung function testing.<sup>19–21</sup> Thirty-one young infants, age range 3–13 weeks, 7 with CF, were studied during behaviorally defined natural sleep. Measurements followed regular feeding of the infants, usually resulting in quiet (non-rapid eye movement) sleep in this age group. Eight older infants with CF aged 13–14 months were studied during chloral-hydrate sedation. Flow was measured using a mainstream ultrasonic flow meter (Exhalyzer D; Eco Medics AG, Duernten, Switzerland).<sup>17</sup> Data were used if no apparent volume drift was present, defined as a change exceeding 2 ml s<sup>-1</sup>. Tidal breathing indices derived from tidal flow were measured from all breaths. Main outcome was V<sub>T</sub> which directly affects end-expiratory lung volume, the functional residual capacity (FRC). Secondary outcomes were respiratory rate (RR), minute ventilation (V<sub>E</sub>), mean tidal expiratory flow (MTEF), and end-tidal CO<sub>2</sub> monitored online. Tidal breathing indices within children were compared by paired *t*-test. *P*-values < 0.05 were considered significant (STATA 11; College Station). Variability of V<sub>T</sub> was estimated by the coefficient of variation (CV = SD/mean %) within each test and per child.

## RESULTS

Thirty-eight infants completed the study according to protocol (Table 1). None of the infants experienced apnea, desaturation (≤92% O<sub>2</sub>), bradycardia (<60 min<sup>-1</sup>) or change in end-tidal CO<sub>2</sub> levels. One non-sedated infant woke up during 100% O<sub>2</sub> exposure (protocol II). In total, V<sub>T</sub>, RR, V<sub>E</sub>, and MTEF were measured in 4,063 breaths from 132 tests.

In both protocols, initial hyperoxia (40% or 100% O<sub>2</sub>) induced hypoventilation. In 33/39 infants, V<sub>T</sub> declined after 2–4 breaths, remained low during 10–25 breaths, and then returned to baseline (Fig. 1). Similarly V<sub>E</sub> and MTEF but not RR decreased and increased again (Table 2). V<sub>T</sub> decreased irrespective of disease, age, or sedation. We stratified children according to disease, age, and sedation: Mean (95% CI) V<sub>T</sub> decrease was 2.9 (1.4; 4.4) ml in the 24 controls, and 6.7 (2.9; 10.4) ml in the 15 infants with CF; 3.0 (1.7; 4.2) ml in the 31 non-sedated (younger) infants, and 9.7 (3.3; 16.2) ml in the eight sedated infants (older, all with CF). Intra-individual variability of V<sub>T</sub> increased during initial hyperoxia (Table 2). In both protocols, mean (SD) CV rose from 8.8 (3.6)% at baseline to 15.1 (7.3)%, *P* < 0.001, during initial hyperoxia. Variability of V<sub>T</sub> increased irrespective of age, sedation, or disease (*P* ≤ 0.001 for all).

The infants' respiratory control apparently adapted to hyperoxia (Figs. 1 and 2). Here, we report ventilatory



**Fig. 1.** Tidal breathing pattern during the classical and new nitrogen MBW protocol. The course of tidal volumes ( $V_T$ ) from 39 infants (16 with CF of which 8 were sedated) across both protocols illustrates transient hypoventilation induced by hyperoxia (40% or 100%  $O_2$ ). The bars give median and inter-quartile ranges of  $V_T$ .  $V_T$  slowly returns to baseline during both 100%  $O_2$  (protocol I: 23 infants, 4 sedated; upper panel) and 40%  $O_2$  exposure (protocol II: 16 infants, 4 sedated; lower panel) and remains stable during subsequent 100%  $O_2$  exposure in protocol II.

dynamics from infants in protocol (II). After initial  $V_T$  decrease during 40%  $O_2$ ,  $V_T$  returned to baseline during 100%  $O_2$  exposure. Mean (SD) baseline  $V_T$  was 51.4 (29.2) ml,  $V_T$  during 40%  $O_2$  exposure was 46.4 (26.8) ml,  $P = 0.007$  and 51.5 (29.2) ml,  $P > 0.9$  during 100%  $O_2$  exposure. Similarly  $V_E$  and MTEF returned to baseline while RR remained unchanged. Similarly end-tidal  $CO_2$  levels monitored online during hyperoxia remained stable.  $V_T$  variability remained slightly elevated (Table 2).

## DISCUSSION

### Summary

This is the first study to report on a new infant  $N_2$ MBW protocol inducing tolerance to hyperoxia. Infant  $N_2$ MBW protocols using 100%  $O_2$  or both 40% and 100%  $O_2$

applied sequentially are safe. Both protocols initially lead to hypoventilation and increased variability of  $V_T$ . These  $V_T$  dynamics were apparently independent of age, disease, and natural or induced sleep in infants younger than 14 months. Of importance, the infants' hypoventilation is temporary. The new protocol comprising 40%  $O_2$  exposure prior to  $N_2$ MBW using 100%  $O_2$  induces tolerance to hyperoxia. Breathing 40%  $O_2$  allows later washout of the lungs' remaining  $N_2$  fraction using pure  $O_2$ . The new protocol seems applicable for future  $N_2$ MBW studies in healthy infants and infants with CF.

### Physiological Aspects of Nitrogen Multiple-Breath Washout in Infants

The principle of  $N_2$ MBW is based on lung inherent  $N_2$  washout via breathing pure  $O_2$ . It appears that temporary

**TABLE 2—Impact of the Classical and New Nitrogen MBW Protocol on Tidal Breathing**

	Baseline	Protocol I	Protocol II	
	Ambient air	100% O <sub>2</sub>	40% O <sub>2</sub>	100% O <sub>2</sub>
N	39	23	16	15
	Mean [range]	Mean diff. (95% CI)	Mean diff. (95% CI)	Mean diff. (95% CI)
V <sub>T</sub> (ml)	44.7 [23.4–121.0]	−3.9 (−6.1; −1.7) <sup>#</sup>	−4.9 (−8.0; −1.9) <sup>#</sup>	0.1 (−2.2; 2.4) <sup>*</sup>
CV <sub>V<sub>T</sub></sub> (%)	8.8 [4.3–23.1]	8.7 (5.9; 11.5) <sup>#</sup>	2.2 (−0.8; 5.1) <sup>*</sup>	2.1 (−0.8; 5.1) <sup>*</sup>
RR (min <sup>−1</sup> )	39 [24–59]	−1 (−3; 1) <sup>*</sup>	1 (−2; 4) <sup>*</sup>	0 (−3; 3) <sup>*</sup>
V <sub>E</sub> (ml min <sup>−1</sup> )	1,575 [980–2,993]	−177 (−246; −108) <sup>#</sup>	−141 (−251; −31) <sup>##</sup>	−9 (−102; 119) <sup>*</sup>
MTEF (ml s <sup>−1</sup> )	46.1 [27.1–80.5]	−6.5 (−8.9; −4.0) <sup>#</sup>	−3.4 (−7.1; 0.3) <sup>*</sup>	0.9 (−2.7; 4.5) <sup>*</sup>

Tidal breathing indices from baseline are given as mean [range], and as mean difference (95% confidence intervals) between (...) baseline and later tests. One infant woke up during breathing 100% O<sub>2</sub>. V<sub>T</sub>, tidal volume; CV<sub>V<sub>T</sub></sub>, intra-subject coefficient of variation from tidal volume; RR, respiratory rate; V<sub>E</sub>, minute ventilation; MTEF, mean tidal expiratory flow. Paired *t*-tests:

<sup>#</sup>*P* ≤ 0.003;

<sup>##</sup>*P* = 0.015;


<sup>\*</sup>*P* ≥ 0.100.

hyperoxia does not influence respiratory control in children older than 4 years,<sup>22</sup> but may trigger the infants' peripheral chemoreceptors which then activate the brainstem respiratory pattern generator network.<sup>15</sup> Hyperoxia may also induce temporary vasoconstriction of cerebral circulation, an effect that theoretically could have contributed to the observed change in ventilation by the induced local increase in tissue CO<sub>2</sub>. Alternative pathways mediated by systemic hypercapnia seem less likely as end-tidal CO<sub>2</sub> levels monitored throughout the studies remained stable. During hyperoxia, the respiratory control adapted. Administration of 40% O<sub>2</sub> induced tolerance to hyperoxia leading to normal but slightly more

variable V<sub>T</sub> during 100% O<sub>2</sub> exposure. Of note, breathing pattern changed in some but not all infants (Fig. 2). Respiratory responses were heterogeneous and apparently independent of age, disease, and sedation.

### Methodological Aspects of Nitrogen Multiple-Breath Washout in Infants

Previous MBW techniques based on sulfur hexafluoride or helium are safe and have been frequently used in children. However, sulfur hexafluoride is a potent greenhouse gas and not available in many countries.<sup>1</sup> Helium is highly volatile which increases the risk of gas



**Fig. 2. Impact of the classical and new nitrogen MBW protocol on tidal volumes.** Mean tidal volumes during nitrogen multiple-breath washout (N<sub>2</sub>MBW) in 23 healthy infants (closed circles) and 16 infants with CF (hollow circles) of which 8 were sedated. In the classical 1-step protocol, infants were exposed to ambient air and 100% oxygen (O<sub>2</sub>). In the new 2-steps protocol, infants were exposed to ambient air, 40% O<sub>2</sub>, and 100% O<sub>2</sub>. Classical protocol on the left panel: 23 infants, 4 sedated; new protocol on the right panel: 16 infants, 4 sedated.

leakage during infant MBW. N<sub>2</sub>MBW has regained interest for lung function testing in infants and young children as N<sub>2</sub>MBW seems appropriate for widespread use. N<sub>2</sub>MBW setups are available and O<sub>2</sub> supply is easily accessible. Modern N<sub>2</sub>MBW setups provide online tracing of expired N<sub>2</sub> and flow-volume loops.<sup>22,23</sup> These technical features are also recommended by the new ERS/ATS consensus statement on inert gas washout testing<sup>1</sup> and could make the new N<sub>2</sub>MBW protocol attractive for early CF lung disease assessment.<sup>12,22,24,25</sup> Previous studies<sup>12,13</sup> reported initial hypoventilation during classical N<sub>2</sub>MBW (100% O<sub>2</sub>) in healthy infants and infants with CF, bronchopulmonary dysplasia, or wheezy bronchitis. However analysis of the breathing pattern was presumably done post hoc as online V<sub>T</sub> control in previous N<sub>2</sub>MBW setups was not available.

### Strengths and Limitations of the Study

The ventilatory response to hyperoxia was similar in older sedated sleeping CF infants and younger non-sedated sleeping CF infants. However, carotid chemoreceptor and ventilatory response to hypoxia are known to undergo significant changes early in life. Due to ethical considerations, we did not sedate younger infants as MBW success rate is excellent without sedation in this age group.<sup>26</sup> We acknowledge that consecutive allocation hampered the assessment of the magnitude of the ventilatory chemoreflex and possible interactions between ventilatory response and age and sedation. Allocation ensured equal distribution of healthy and diseased infants and older infants requiring sedation in both protocols.

The current N<sub>2</sub>MBW setup is accurate for measuring infant lung volumes as recently shown in a plastic lung model.<sup>23</sup> Due to expected software adaptations for 40% O<sub>2</sub>, we could not measure classical in vivo N<sub>2</sub>MBW outcomes such as the LCI which may limit the interpretation of the current data. However, because of the complex dynamic respiratory control and the likely change in FRC, changes in, for example, LCI seem also likely (LCI = CEV/FRC).<sup>13</sup> Furthermore, the observed initial depression of the first few breaths when hyperoxia is administered is of importance. The depression in ventilation affects our potential to derive alternative indices of gas mixing efficiency based on phase III slopes (Sacin and Scand). The latter assumptions are supported by recent preliminary data from school-aged children suggesting a significant decrease in gas mixing efficiency upon change in breathing pattern.<sup>27</sup> The impact from both N<sub>2</sub>MBW protocols on FRC and LCI in infants nevertheless remains to be determined. Before this is done, MBW tests using foreign inert tracer gases seem preferable in infants.

### CONCLUSION

Taken together, we provide new perspectives for future N<sub>2</sub>MBW infant studies. Hyperoxia for the purpose of pulmonary function testing is safe but influences ventilatory control in infants. The new 40/100% O<sub>2</sub> protocol comprising 40% O<sub>2</sub> exposure prior to N<sub>2</sub>MBW using 100% O<sub>2</sub> may induce tolerance to hyperoxia. The current technique requires adaption for gradual O<sub>2</sub> administration in infants. We assume that continued 100% O<sub>2</sub> exposure, for example for 60 breaths, would similarly induce tolerance to hyperoxia but would not allow subsequent N<sub>2</sub>MBW. The new 40/100% O<sub>2</sub> protocol seems an attractive option for N<sub>2</sub>MBW in infants.

### ACKNOWLEDGMENTS

The authors thank all children and their families for their participation in the study. The authors would like to express their gratitude to A. Lindblad (Queen Silvia Children's Hospital, Gothenburg, Sweden), F. Ratjen (Hospital for Sick Children, Toronto, Canada), and T. Riedel (Department of Paediatrics, University Hospital of Bern, Switzerland) for their valuable input. The authors thank G. Cangiano, S. Luescher, M. Graf, G. Wirz, and S. Schmid (Department of Paediatrics, University Hospital of Bern, Switzerland) for their valuable work and support. We acknowledge the valuable work of the "Swiss Cystic Fibrosis Newborn Screening Group": Constance Barazzone, Matthias R. Baumgartner, Anne Mornand, Peter Eng, Ralph Fingerhut, Sabina Gallati, Gaudenz Hafen, Jürg Hammer, Claudia E. Kühni, Alexander Möller, Dominik Müller, Isabelle Rochat, Corina S. Rueegg, Barbara Schiller, Martin H. Schöni, Renate Spinaz, Johannes Spalinger, Daniel Trachsel, Toni Torresani, and Maura Zanolari (ordered alphabetically).

### REFERENCES

1. Robinson PD, Latzin P, Verbanck S, Hall GL, Horsley A, Gappa M, Thamrin C, Arets HG, Aurora P, Fuchs SI, King GG, Lum S, Macleod K, Paiva M, Pillow JJ, Ranganathan S, Ratjen F, Singer F, Sonnappa S, Stocks J, Subbarao P, Thompson BR, Gustafsson PM. Consensus statement for inert gas washout measurement using multiple- and single-breath tests. *Eur Respir J* 2013;41:507–522.
2. Kieninger E, Singer F, Fuchs O, Abbas C, Frey U, Regamey N, Casaulta C, Latzin P. Long-term course of lung clearance index between infancy and school-age in cystic fibrosis subjects. *J Cyst Fibros* 2011;10:487–490.
3. Gustafsson PM, Aurora P, Lindblad A. Evaluation of ventilation maldistribution as an early indicator of lung disease in children with cystic fibrosis. *Eur Respir J* 2003;22:972–979.
4. Fuchs SI, Toussaint S, Edlhaime B, Ballmann M, Gappa M. Short-term effect of physiotherapy on variability of the lung clearance index in children with cystic fibrosis. *Pediatr Pulmonol* 2010;45:301–306.

5. Gustafsson PM, De Jong PA, Tiddens HA, Lindblad A. Multiple-breath inert gas washout and spirometry versus structural lung disease in cystic fibrosis. *Thorax* 2008;63:129–134.
6. Owens CM, Aurora P, Stanojevic S, Bush A, Wade A, Oliver C, Calder A, Price J, Carr SB, Shankar A, Stocks J. Lung Clearance Index and HRCT are complementary markers of lung abnormalities in young children with CF. *Thorax* 2011;66:481–488.
7. Hall GL, Logie KM, Parsons F, Schulzke SM, Nolan G, Murray C, Ranganathan S, Robinson P, Sly PD, Stick SM. Air trapping on chest CT is associated with worse ventilation distribution in infants with cystic fibrosis diagnosed following newborn screening. *PLoS ONE* 2011;6:e23932.
8. Aurora P, Stanojevic S, Wade A, Oliver C, Kozłowska W, Lum S, Bush A, Price J, Carr SB, Shankar A, Stocks J. Lung clearance index at 4 years predicts subsequent lung function in children with cystic fibrosis. *Am J Respir Crit Care Med* 2011;183:752–758.
9. Amin R, Subbarao P, Lou W, Jabar A, Balkovec S, Jensen R, Kerrigan S, Gustafsson P, Ratjen F. The effect of dornase alfa on ventilation inhomogeneity in patients with Cystic Fibrosis. *Eur Respir J* 2011;37:806–812.
10. Amin R, Subbarao P, Jabar A, Balkovec S, Jensen R, Kerrigan S, Gustafsson P, Ratjen F. Hypertonic saline improves the LCI in paediatric patients with CF with normal lung function. *Thorax* 2010;65:379–383.
11. Horsley AR, Davies JC, Gray RD, Macleod KA, Donovan J, Aziz ZA, Bell NJ, Rainer M, Mt-Isa S, Voase N, Dewar MH, Saunders C, Gibson JS, Parra-Leiton J, Larsen MD, Jeswiet S, Soussi S, Bakar Y, Meister MG, Tyler P, Doherty A, Hansell DM, Ashby D, Hyde SC, Gill DR, Greening AP, Porteous DJ, Innes JA, Boyd AC, Griesenbach U, Cunningham S, Alton EW. Changes in physiological, functional and structural markers of cystic fibrosis lung disease with treatment of a pulmonary exacerbation. *Thorax* 2013;68:532–539.
12. Schibler A, Schneider M, Frey U, Kraemer R. Moment ratio analysis of multiple breath nitrogen washout in infants with lung disease. *Eur Respir J* 2000;15:1094–1101.
13. Gustafsson PM, Källman S, Bhiladvala M, Fletcher ME. N<sub>2</sub>-washout with pure oxygen reduces tidal volumes markedly in sedated infants. *Eur Respir J* 1997;10(Suppl. 25):165s.
14. Hertzberg T, Lagercrantz H. Postnatal sensitivity of the peripheral chemoreceptors in newborn infants. *Arch Dis Child* 1987;62:1238–1241.
15. Chardon K, Telliez F, Bach V, Leke A, Delanaud S, Bouferrache B, Libert JP, Gaultier C. Effects of warm and cool thermal conditions on ventilatory responses to hyperoxic test in neonates. *Respir Physiol Neurobiol* 2004;140:145–153.
16. Fuchs O, Latzin P, Kuehni CE, Frey U. Cohort profile: the Bern infant lung development cohort. *Int J Epidemiol* 2012;41:366–376.
17. Fuchs O, Latzin P, Thamrin C, Stern G, Frischknecht P, Singer F, Kieninger E, Proietti E, Riedel T, Frey U. Normative data for lung function and exhaled nitric oxide in unsedated healthy infants. *Eur Respir J* 2011;37:1208–1216.
18. Barben J, Gallati S, Fingerhut R, Schoeni MH, Baumgartner MR, Torresani T. Retrospective analysis of stored dried blood spots from children with cystic fibrosis and matched controls to assess the performance of a proposed newborn screening protocol in Switzerland. *J Cyst Fibros* 2012;11:332–336.
19. Frey U, Stocks J, Coates A, Sly P, Bates J. Specifications for equipment used for infant pulmonary function testing. ERS/ATS Task Force on Standards for Infant Respiratory Function Testing. European Respiratory Society/American Thoracic Society. *Eur Respir J* 2000;16:731–740.
20. Frey U, Stocks J, Sly P, Bates J. Specification for signal processing and data handling used for infant pulmonary function testing. ERS/ATS Task Force on Standards for Infant Respiratory Function Testing. European Respiratory Society/American Thoracic Society. *Eur Respir J* 2000;16:1016–1022.
21. Bates JH, Schmalisch G, Filbrun D, Stocks J. Tidal breath analysis for infant pulmonary function testing. ERS/ATS Task Force on Standards for Infant Respiratory Function Testing. European Respiratory Society/American Thoracic Society. *Eur Respir J* 2000;16:1180–1192.
22. Singer F, Kieninger E, Abbas C, Yammine S, Fuchs O, Proietti E, Regamey N, Casaulta C, Frey U, Latzin P. Practicability of nitrogen multiple-breath washout measurements in a pediatric cystic fibrosis outpatient setting. *Pediatr Pulmonol* 2012; doi: 10.1002/ppul.22651
23. Singer F, Houlzt B, Latzin P, Robinson P, Gustafsson P. A realistic validation study of a new nitrogen multiple-breath washout system. *PLoS ONE* 2012;7:e36083.
24. Verbanck S, Paiva M, Paeps E, Schuermans D, Malfroot A, Vincken W, Vanderhelst E. Lung clearance index in adult CF patients: the role of convection-dependent lung units. *Eur Respir J* 2012; doi: 10.1183/09031936.00125312
25. Gappa M, Fletcher ME, Dezateux CA, Stocks J. Comparison of nitrogen washout and plethysmographic measurements of lung volume in healthy infants. *Am Rev Respir Dis* 1993;148:1496–1501.
26. Latzin P, Roth S, Thamrin C, Hutten GJ, Pramana I, Kuehni CE, Casaulta C, Nelle M, Riedel T, Frey U. Lung volume, breathing pattern and ventilation inhomogeneity in preterm and term infants. *PLoS ONE* 2009;4:e4635.
27. Yammine S, Singer F, Latzin P. One size does not fit all—impact of the one liter tidal volume breathing protocol on indices from nitrogen multiple-breath washout in children. *Eur Respir J* 2013; 40:259s.